

Feynman’s clock, a new variational principle, and quantum time-orbitals

Jarrold R. McClean,¹ John A. Parkhill,¹ and Alán Aspuru-Guzik¹

¹*Department of Chemistry and Chemical Biology, Harvard University, Cambridge MA, 02138*

We introduce a new discrete-time variational principle inspired by the quantum clock originally proposed by Feynman, and use it to write down quantum evolution as a ground state eigenvalue problem. This allows one to apply ground state quantum many-body theory to quantum dynamics, extending the reach of many highly developed tools from this fertile research area. We draw an explicit connection between previously known time-dependent variational principles and the new time embedded variational principle presented. Sample calculations are presented applying the idea to the electronic degrees of freedom in a Hydrogen molecule and the spin degrees of freedom of a model inorganic compound. We employ the configuration-interaction approximation on a basis in which time is a variable. Finally, we take advantage of the unique perspective of the new variational principle to examine the error of basis approximations in quantum dynamics.

Introduction

Feynman proposed a revolutionary solution to the problem of quantum simulation: use quantum computers to simulate quantum systems. While this strategy is powerful and elegant, universal quantum computers may not be available for some time, and in fact, accurate quantum simulations may be required for their eventual construction. In this work, we will use the clock Hamiltonian originally introduced by Feynman [1, 2] for the purposes of quantum computation, to re-write the quantum dynamics problem as a ground state eigenvalue problem. We then generalize this approach, and obtain a novel variational principle for the dynamics of a quantum system. Variational principles play a central role in the development and study of quantum dynamics [3–10], and the variational principle presented here extends the arsenal of available tools by allowing one to directly apply efficient approximations from the ground state quantum many-body problem to study dynamics.

Previous approaches for recasting quantum dynamics as a time-independent problem include Floquet theory for periodic potentials [11–13] and more generally the (t, t') formalism of Peskin and Moiseyev [14]. However, the approach proposed in this manuscript differs considerably from these previous approaches. We derive our result from a different variational principle, and in our embedding the dynamics of the problem are encoded directly in its solution, as opposed to requiring the construction of another propagator. Moreover, the method of solution we describe will differ considerably from previous approaches by employing techniques from ground state many-body theory to the time-dependent path as a whole. Considerable work has now been done in the migration of knowledge from classical computing to quantum computing and quantum information [15]. In this paper, we propose a novel use of an idea from quantum computation for the simulation of quantum dynamics.

The successful application of quantum computers to quantum chemistry [16, 17] has been demonstrated, in

both theory [18–21] and experiment [22, 23]. The exact classical simulation of quantum dynamics is a challenging problem due to the exponentially large space in which the problem must be solved. To make progress using classical computers, accurate approximate schemes with controllable errors must be developed. Some ideas from quantum information have successfully led to new methods for simulating many-body states on classical computers [24, 25]. More generally, quantum information tools [15, 26] have become widely applicable in a variety of settings, such as understanding the role of entanglement in many-body systems [27].

The paper is organized as follows. We will first review the Feynman clock: a mapping stemming from the field of quantum computation that can be employed for converting problems in quantum evolution into ground-state problems in a larger Hilbert space. We then generalize the Feynman clock into a time-embedded discrete variational principle (TEDVP) which offers additional insight to quantum time-dynamics in a way that is complementary to existing differential variational principles [28–30]. The basic dynamics of an excited Hydrogen molecule are used to explain how dynamics is simulated with the clock Hamiltonian. We then apply the configuration interaction method [31, 32] from quantum chemistry to solve for approximate dynamics of a model spin system demonstrating convergence of accuracy of our proposed approach with level of the truncation. Finally we discuss metrics inspired by our approach that can be used to quantitatively understand the errors resulting from truncating the Hilbert space of many-body quantum dynamics.

Physical dynamics as a sequence of quantum gates

Consider a quantum system described by a time-dependent wavefunction $|\Psi(t)\rangle$. The dynamics of this system are determined by a Hermitian Hamiltonian $H(t)$ according to the time-dependent Schrödinger equation in

atomic units,

$$i\partial_t |\Psi(t)\rangle = H(t) |\Psi(t)\rangle \quad (1)$$

A formal solution to the above equation can be generally written:

$$|\Psi(t)\rangle = \mathcal{T} \left(e^{-i \int_{t_0}^t dt' H(t')} \right) |\Psi(t_0)\rangle = U(t, t_0) |\Psi(t_0)\rangle \quad (2)$$

Where \mathcal{T} is the well known time-ordering operator and $U(t, t_0)$ is a unitary operator that evolves the system from a time t_0 to a time t . These operators also obey a group composition property, such that if $t_0 < t_1 < \dots < t_n < t$ then

$$|\Psi(t)\rangle = U(t, t_0) |\Psi(t_0)\rangle = U(t, t_n) U(t_n, t_{n-1}) \dots U(t_1, t_0) |\Psi(t_0)\rangle \quad (3)$$

From the unitarity of these operators, it is of course also true that $U(t_n, t_{n-1})^\dagger = U(t_{n-1}, t_n)$ where \dagger indicates the adjoint. Thus far, we have treated time as a continuous variable. However, when one considers numerical calculations on a wavefunction, it is typically necessary to discretize time.

We discretize time by keeping an ancillary quantum system, which can occupy states with integer indices ranging from $|0\rangle$ to $|T-1\rangle$ where T is the number of discrete time steps under consideration. This quantum system has orthonormal states such that

$$\langle i | j \rangle = \delta_{ij} \quad (4)$$

This definition allows one to encode the entire evolution of a physical system by entangling the physical wavefunction with this auxiliary quantum system representing time, known as the “time register”. We define this compound state to be the history state, given by

$$|\Phi\rangle = \frac{1}{\sqrt{T}} \sum_t |\Psi_t\rangle \otimes |t\rangle \quad (5)$$

where subscripts are used to emphasize when we are considering a time-independent state of a system at time t . That is, we define $|\Psi_i\rangle = |\Psi(t)\rangle|_{t=t_i}$. From these definitions, it is immediately clear from above that the wavefunction at any time t can be recovered by projection with the time register, such that

$$|\Psi(t)\rangle|_{t=t_i} = \sqrt{T} \langle i | \Phi \rangle \quad (6)$$

Additionally, we discretize the action of our unitary operators, such that $U(t_1, t_0) = U_0$ and we embed this operator into the composite system-time Hilbert space as $(U_0 \otimes |1\rangle \langle 0|)$. While this notation is standard in the field of quantum information, it is less prevalent in chemical physics. For clarity, we offer an example algebraic manipulation in the supplementary information.

While the utility of this discretization has not yet been made apparent, we will now use this discretization to transform the quantum dynamics problem into a ground state eigenvalue problem.




FIG. 1. A schematic representation of the action of the clock Hamiltonian on the history state with three discrete times, and a Hilbert space of three states. Each block is a matrix with dimension of the physical system.

Feynman’s clock

In the gate model [15, 33] of quantum computation, one begins with an initial quantum state $|\Psi_0\rangle$ state, applies a sequence of unitary operators $\{U_i\}$, known as quantum gates. By making a measurement on the final state $|\Psi_f\rangle$, one determines the result of the computation, or equivalently the result of applying the sequence of unitary operators $\{U_i\}$. The map from the sequence of unitary operators $\{U_i\}$ in the gate model, to a Hamiltonian that has the clock state as its lowest eigenvector is given by a construction called the Clock Hamiltonian [2]. Since its initial inception, much work has been done on the specific form of the clock, making it amenable to implementation on quantum computers [34]. However, for the purposes of our discussion that pertain to implementation on a classical computer, the following simple construction suffices

$$\mathcal{H} = C_0 + \frac{1}{2} \sum_{t=0}^{T-1} (I \otimes |t\rangle \langle t| - U_t \otimes |t+1\rangle \langle t| - U_t^\dagger \otimes |t\rangle \langle t+1| + I \otimes |t+1\rangle \langle t+1|) \quad (7)$$

where C_0 is a penalty term which can be used to specify the state of the physical system at any time. Typically, we use this to enforce the initial state, such that if the known state at time $t = 0$ is given by $|\Psi_0\rangle$. For this, we define:

$$C_0 = (I - |\Psi_0\rangle \langle \Psi_0|) \otimes |0\rangle \langle 0| \quad (8)$$

One may verify by action of \mathcal{H} on the history state defined above, $|\Phi\rangle$, that the history state is an eigenvector of this operator, with eigenvalue 0.

A discrete-time variational principle

We now introduce a new time-embedded discrete variational principle (TEDVP) and show how the above eigenvalue problem emerges as the special case of choosing a linear variational space. Suppose that one knows the exact form of the evolution operators and wavefunctions at times $t = 0, 1$. By the properties of unitary evolution it is clear that the following holds:

$$2 - \langle \Psi_1 | U_0 | \Psi_0 \rangle - \langle \Psi_0 | U_0^\dagger | \Psi_1 \rangle = 0 \quad (9)$$

From this, we can see that if the exact construction of U_i is known for all i , but the wavefunctions are only approximately known (but still normalized), it is true that

$$\sum_{t=0}^{T-1} \left(2 - \langle \Psi_{t+1} | U_t | \Psi_t \rangle - \langle \Psi_t | U_t^\dagger | \Psi_{t+1} \rangle \right) \geq 0 \quad (10)$$

where equality holds in the case that the wavefunction becomes exact at each discrete time point. Reintroducing the ancillary time-register, we may equivalently say that all valid time evolutions embedded into the composite system-time space as $\sum_t |\Psi_t\rangle |t\rangle$ minimize the quantity

$$\mathcal{S} = \sum_{t,t'} \langle t' | \langle \Psi_{t'} | \mathcal{H} | \Psi_t \rangle | t \rangle \quad (11)$$

where \mathcal{H} (script font for operators denotes they act in the composite system-time space) is the operator given by

$$\begin{aligned} \mathcal{H} = & \sum_{t=0}^{T-1} I \otimes |t\rangle \langle t| - U_t \otimes |t+1\rangle \langle t| \\ & - U_t^\dagger \otimes |t\rangle \langle t+1| + I \otimes |t+1\rangle \langle t+1| \end{aligned} \quad (12)$$

It is then clear from the usual ground state variational principle, that the expectation value of the operator

$$\mathcal{S} = \sum_{t,t'} \langle t' | \langle \Psi_{t'} | \mathcal{H} | \Psi_t \rangle | t \rangle \quad (13)$$

is only minimized for exact evolutions of the physical system. This leads us immediately to a time-dependent variational principle for the discrete representation of a wavefunction given by:

$$\delta \mathcal{S} = \delta \sum_{t,t'} \langle t' | \langle \Psi_{t'} | \mathcal{H} | \Psi_t \rangle | t \rangle = 0 \quad (14)$$

It is interesting to note, that this is a true variational principle in the sense that an exact quantum evolution is found at a minimum, rather than a stationary point as in some variational principles [30]. This is related to the connection between this variational principle and the McLachlan variational principle that is explored in the

next section. However, to the authors knowledge, this connection has never been explicitly made until now.

To complete the connection to the clock mapping given above, we first note that this operator is Hermitian by construction and choose a linear variational space that spans the entire physical domain. To constrain the solution to have unit norm, we introduce the Lagrange multiplier λ and minimize the auxiliary functional given by

$$\mathcal{L} = \sum_{t,t'} \langle t' | \langle \Psi_{t'} | \mathcal{H} | \Psi_t \rangle | t \rangle + C_0 - \lambda \left(\sum_{t,t'} \langle t' | \langle \Psi_{t'} | \Psi_t \rangle | t \rangle - 1 \right) \quad (15)$$

It is clear that this problem is equivalent to the exact eigenvalue problem on \mathcal{H} with eigenvalue λ . The optimal trajectory is given by the ground state eigenvector of the operator \mathcal{H} . From this construction, we see that the clock mapping originally proposed by Feynman is easily recovered as the optimal variation of the TEDVP in a complete linear basis, under the constraint of unit norm. Note that the inclusion of C_0 as a penalty term to break the degeneracy of the ground state is only a convenient construction for the eigenvalue problem. In the general TEDVP, one need not include this penalty explicitly, as degenerate allowed paths are excluded, as in other time-dependent variational principles, by fixing the initial state.

We note, as in the case of the time-independent variational principle and differential formulations of the time-dependent variational principle, the most compact solutions may be derived from variational spaces that have non-linear parameterizations. Key examples of this in chemistry include Hartree-Fock, coupled cluster theory [35–37], and multi-configurational time-dependent Hartree [38]. It is here that the generality of this new variational principle allows one to explore new solutions to the dynamics of the path without the restriction of writing the problem as an eigenvalue problem, as in the original clock construction of Feynman.

Connection to previous time-dependent variational principles

In the limit of an exact solution, it must be true that all valid time-dependent variational principles are satisfied. For that reason, it is important to draw the formal connection between our variational principle and previously known variational principles.

Considering only two adjacent times t and $t+1$, the operator \mathcal{H} is given by:

$$\begin{aligned} \mathcal{H} = & \frac{1}{2} (I \otimes |t\rangle \langle t| - U_t \otimes |t+1\rangle \langle t| \\ & - U_t^\dagger \otimes |t\rangle \langle t+1| + I \otimes |t+1\rangle \langle t+1|) \end{aligned} \quad (16)$$

Now suppose that the separation of physical times between discrete step t and $t+1$, denoted dt is small, and




FIG. 2. The spin triangle within the vanadium compound, used as a model system for the TEDVP. Note that coordinating sodium ions and water molecules are not depicted here. The chemical formula of this compound is given by $(\text{CN}_3\text{H}_6)_4\text{Na}_2[\text{H}_4\text{V}_6^{(IV)}\text{O}_8(\text{PO}_4)_4(\text{OCH}_2)_3\text{CCH}_2\text{OH}_2] \cdot 14\text{H}_2\text{O}$

the physical system has an associated Hamiltonian given by H , such that

$$U_t = e^{-iHdt} \approx I - iHdt - \frac{H^2 dt^2}{2} \quad (17)$$

By inserting this propagator into equation 14, rewriting the result in terms of $|\Psi(t)\rangle$, and dropping terms of order dt^3 we have

$$\begin{aligned} & \delta (\langle \Psi(t) | \Psi(t) \rangle - \langle \Psi(t+dt) | (I - iHdt) | \Psi(t) \rangle \\ & - \langle \Psi(t) | (I + iHdt) | \Psi(t+dt) \rangle + \\ & \langle \Psi(t) | H^2 dt^2 | \Psi(t) \rangle + \langle \Psi(t+dt) | \Psi(t+dt) \rangle) = 0 \end{aligned} \quad (18)$$

After defining the difference operator such that $\partial_t |\Psi(t)\rangle \equiv [|\Psi(t+dt)\rangle - |\Psi(t)\rangle] / dt$, we can factorize the above expression into

$$\delta \langle \Psi(t) | (i\partial_t - H)^\dagger (i\partial_t - H) | \Psi(t) \rangle = 0 \quad (19)$$

In the limit that $dt \rightarrow 0$, these difference operators become derivatives. Defining $\Theta = i\partial_t |\Psi(t)\rangle$, this is precisely the McLachlan variational principle [30].

$$\delta \|\Theta - H |\Psi(t)\rangle\|^2 = 0 \quad (20)$$

We then conclude that in the limit of infinitesimal physical time for a single time step, the TEDVP is equivalent to the McLachlan variational principle. Under the reasonable assumptions that the variational spaces of the wavefunction and its time derivative are the same and that the parameters are complex analytic, then it is also equivalent to the Dirac-Frenkel and Lagrange variational principles [39]. Moreover, in an appendix, we provide a connection that allows other variational principles to be written as eigenvalue problems, and further discuss the merits of the integrated formalism used here.

To conclude this section, we highlight one additional difference between practical uses of the TEDVP and

other variational principles: The TEDVP is independent of the method used to construct the operator U_t . In quantum information applications, this implies it is not required to know a set of generating Hamiltonians for quantum gates. Additionally, in numerical applications, one is not restricted by the choice of approximate propagator used. In cases where an analytic propagator is known for the chosen basis, it can be sampled explicitly.

Structure of the TEDVP eigenvalue problem

Suppose that the dimension of the physical system is given by N and the number of timesteps of interest is given by T . Assuming that the time register $|T\rangle$ is ordered, the resulting eigenvalue problem is block tridiagonal with dimension NT (See Fig. 1).

This structure has been described at least once before in the context of ground-state quantum computation [40], but to the authors knowledge, never in the context of conventional simulation of quantum systems. The block tridiagonal structure is particularly amenable to the use of parallel sparse eigensolvers. This opens up the possibility of exploiting the TEDVP to carry out simulations of time dynamics in parallel using modern computer architectures. Parallel-in-time physical dynamics [41, 42] has been seldom explored, but will become increasingly important as modern computer core speeds level out, and massive parallelism becomes the only way to accelerate computation.

The solution of eigenvalue problems by sparse linear algebra methods is often limited by the gap between the lowest eigenvalue and the next largest eigenvalue. The spectral properties of the clock Hamiltonian have previously been explored in ground-state quantum computation [43]. Interestingly, it turns out that this Hamiltonian can be viewed as an analog to the tight-binding quantum particle in a box, and has an eigenvalue gap that scales as $O(1/T^2)$, independent of both N and the physical time that the evolution of each operator U_t represents. As the gap between the lowest and next lowest eigenvalue is a determining factor in the difficulty of applying iterative eigenvalue solvers, it is expected that the computational cost of this increase accordingly with the number of time steps taken, independent of the physical time each step represents. It is for this reason that we do not formulate a continuous generalization of our variational principle as in principle the number of time steps in this case tends to infinity. While the authors do not discount the formal value of a continuum construction, the computational tractability of its practical implementations remain unclear.

The eigenvectors of the clock Hamiltonian, which we will denote as time-orbitals, are interesting objects which merit study in their own right. The ground state time-orbital contains the physical dynamics of the system and




FIG. 3. The electron density of the first few clock eigenstates in an H_2 molecule are shown at the 5 percent contour. The lowest state represents the density of the exact correlated dynamics, a slightly left-polarized H_2 evolves for a half a period during the dynamics. Higher states show the localization into short spans of time in non-norm-conserving trajectories. Lobes pinched between the nuclei (black dots) in states 2 and 9 reflect significant double excitation character. With the applied penalty these states differ from one another by less than 3 milliHartrees.

the excited time-orbitals contain orthogonal information. In order to examine the properties of these time-orbitals, we have diagonalized the clock representation of the full configuration interaction of the hydrogen molecule in a minimal basis (STO-3G [44]) beginning from an even superposition of the ground and first excited state. The electron densities of the resulting time orbitals are shown in Fig. 3. The lowest energy time-orbital, which provides the exact quantum dynamics given the initial condition conserves electron density. Interestingly the higher lying states do not conserve probability over simulation time; instead probability localizes in time. The physical meaning of these states, and their time-reversal symmetry breaking is unclear and warrants further study.

TEDVP for approximate dynamics

There has been a recent rise in the interest of methods for simulating quantum spin dynamics in chemistry [45, 46]. To study the properties of the clock mapping when used to formulate approximate dynamics, we chose a simple model spin system inside an inorganic molecule [47]. Specifically, we examine the spin dynamics of the vanadium compound depicted in Fig. 2.

By choosing the three unpaired electron spins to interact with one another by means of isotropic exchange as well as uniform static external magnetic field B_0 , and a time-dependent transverse field B_1 , this system can be modeled with a spin Hamiltonian

$$H = J_a(S_1 \cdot S_2 + S_1 \cdot S_3) + J_c S_2 \cdot S_3 + \mu B_0(S_1^z + S_2^z + S_3^z) + \mu B_1(S_1^x + S_2^x + S_3^x) \quad (21)$$

Where S_i^α is the α Pauli operator on spin i , $\mu = g\mu_b$, μ_b is the Bohr magneton, g is the measured spectroscopic

splitting factor. The couplings $J_a \neq J_c$ as well as g are fitted through experimental determinations of magnetic susceptibility [47]. The fact that they are not equal is reflective of the isosceles geometry of the vanadium centers. The parameters of this model, are given by: $g = 1.95$, $J_a = 64.6 \pm 0.5 K$, and $J_c = 6.9 \pm 1 K$. We will allow B_0 to vary to study the properties of the clock mapping in the solution of approximate quantum dynamics.

Dynamics in a linear basis

The quantum chemistry community has decades of experience in developing and employing methods for obtaining approximate solutions of high-dimensional, ground-state eigenvector problems. By utilizing the connection we have made from dynamics to ground state problem, we will now borrow and apply the most conceptually simple approximation from quantum chemistry: configuration interaction in the space of trajectories [48], to our model problem to elucidate the properties of the clock mapping.

For the uncorrelated reference, we take the entire path of a mean-field spin evolution that is governed by the time-dependent Hartree equations, and write it as:

$$|\Psi_{MF}\rangle = \sum_t \left(\prod_i U_t^i |0\rangle_i \right) |t\rangle \quad (22)$$

$$= \sum_t \left(\prod_i |0\rangle_i^t \right) |t\rangle \quad (23)$$

$$= \sum_t |\phi_t\rangle |t\rangle \quad (24)$$

where $|0\rangle_i$ is the reference spin-down state for spin i , $|0\rangle_i^t$ is the reference spin-down state after rotation at time t ,




FIG. 4. The convergence of the CI expansion *in time* is robust to strength of perturbation, and choice of initial state for the Vanadium complex. In these plots the error of the expectation value of S_z at site 1 is plotted as a function of time, and coupling constant, each propagation with a different coupling constant is begun from a different random initial product state.

$|\phi_t\rangle$ is the whole product system at time t , and U_t^i is determined from the mean field Hamiltonian. The equations of motion that determine U_t^i are derived in a manner analogous to the time-dependent Hartree equations, and if one writes the wavefunction in the physical space as

$$|\psi(t)\rangle = a(t) \prod_i U_i(t) |0\rangle_i = a(t) \prod_i |\phi_i\rangle = a(t) |\Phi(t)\rangle \quad (25)$$

Then the equations of motion are given by

$$a(t) = a(0) \quad (26)$$

$$i\dot{U}_i = (H^{(i)} - \left(\frac{f-1}{f}\right) E(t)) U_i \quad (27)$$

Where $H^{(i)}$ is the mean field Hamiltonian for spin i formed by contracting the Hamiltonian over all other spins $j \neq i$, $E(t)$ is the expectation value of the Hamiltonian at time t , and f is the number of spins in the system.

To generate configurations, we also introduce the transformed spin excitation operator \tilde{S}_{it}^+ , which is defined by

$$S_i^+ |0\rangle_i = |1\rangle_i \quad (28)$$

$$U_t^{i\dagger} S_i^+ U_t^i = \tilde{S}_{it}^+ \quad (29)$$

$$\tilde{S}_{it}^+ |0\rangle_i^t = |1\rangle_i^t \quad (30)$$

In analogy to traditional configuration interaction, we will define different levels of excitation (e.g. singles, doubles, ...) as the number of spin excitations at each time t that will be included in the basis in which the problem is diagonalized. For example, the basis for the configuration interaction singles (CIS) problem is defined as

$$\mathcal{B}_{CIS} = \left\{ \tilde{S}_{jt}^+ |\phi_t\rangle |t\rangle \mid j \in [0, n], t \in [0, T] \right\} \quad (31)$$

Note that \tilde{S}_{jt}^+ for $j = n$ is simply defined to be the identity operator on all spins so that the reference configuration is naturally included. Similarly, the basis for single and double excitations (CISD) is given by

$$\mathcal{B}_{CISD} = \left\{ \tilde{S}_{jt}^+ \tilde{S}_{kt}^+ |\phi_t\rangle |t\rangle \mid j \in [0, n], k \in [0, j], t \in [0, T] \right\} \quad (32)$$

Higher levels of excitation follow naturally, and it is clear that when one reaches a level of excitation equivalent to the number of spins, this method may be regarded as full configuration interaction, or exact diagonalization in the space of discretized paths.

The choice of a time-dependent reference allows the reference configuration to be nearly exact when $B_0, B_1 \gg J_a, J_c$, independent of the nature of the time-dependent transverse field. This allows for the separation of the consequences of time-dependence from the effects of two versus one body spin interactions.

After a choice of orthonormal basis, the dynamics of the physical system are given by the ground state eigenvector of the projected eigenvalue problem

$$\mathcal{H}_{\mathcal{B}_i} |\tilde{\Phi}\rangle = E |\tilde{\Phi}\rangle \quad (33)$$

where we explicitly define the projection operator onto the basis \mathcal{B}_i as $P_{\mathcal{B}_i} = \sum_{|j\rangle \in \mathcal{B}_i} |j\rangle \langle j|$ so that the projected operator is given by $\mathcal{H}_{\mathcal{B}_i} = (P_{\mathcal{B}_i} \mathcal{H} P_{\mathcal{B}_i})$

Using these constructions, we calculate the quantum dynamics of the sample system described above. For convenience, we rescale the Hamiltonian by the value of $1/\mu B_0$. To add arbitrary non-trivial time dependence to the system and mimic the interaction of the system with a transverse pulse, we take $B_1 \propto \exp(-t^2/2) \cos(mt)$. The magnitude of B_0 was taken to be $200T$ in order to model perturbative two body interactions in this Hamiltonian. To propagate the equations of motion and generate the propagators for the clock operator we use the




FIG. 5. The total contribution to the ground state eigenvector from each level of excitation is plotted, where 0=MF, 1=CIS, 2=CISD, and 3=FCI(Exact), and seen to decrease with excitation supporting the quality of the time-dependent reference state. The deviation of the wavefunction norm from unity resulting from projection is seen to decrease monotonically with level of excitation.

enforced time-reversal symmetry exponential propagator [49] given by

$$U_t = \exp\left(-i\frac{dt}{2}H(t+dt)\right)\exp\left(-i\frac{dt}{2}H(t)\right) \quad (34)$$

The dynamics of some physical observables are displayed (Fig. 6) for the reference configuration, single excitations, double excitations, and full configuration interaction. The physical observables have been calculated with normalization at each time step. It is seen (Fig. 4) that as in the case of ground state electronic structure the physical observables become more accurate both qualitatively and quantitatively as the configuration space expands, converging to the exact solution with full configuration interaction. Moreover in Fig. 5 we plot the contributions from the reference configuration, singles space, doubles space, and triples space and observe rapidly diminishing contributions. This suggests that the time-dependent path reference used here provides a good qualitative description of the system. As a result, perturbative and other dynamically correlated methods from quantum chemistry may also be amenable to the solution of this problem.




FIG. 6. Trajectories for single-particle observables in different levels of basis set truncation starting from the reference configuration path (MF). As more levels of spin excitation are included with configuration interaction singles(CIS) and configuration interaction with singles and doubles(CISD) the trajectories converge to the exact result.

In principle, approximate dynamics derived from this variational principle are not norm conserving, as is seen in Fig. 5, however this actually offers an important insight into a major source of error in quantum dynamics simulations of many-body systems, which is the truncation of the basis set as described in the following section. Simulation based on the conventional variational principles that propagate within an incomplete configuration space easily preserve norm; however the trajectories of probability which should have left the simulated space are necessarily in error.

Norm Loss as a measure of truncation error

Conservation of the norm of a wavefunction is often considered a critical property for approximate quantum dynamics, as it is a property of the exact propagator resulting from time-reversal symmetry. However, if norm conservation is enforced in a propagation which does not span the complete Hilbert-space, one must account for the components of the wavefunction that would have evolved into the space not included in the computation. It's not immediately clear how population density which attempts to leave the selected subspace should be folded back into the system without being able to simulate the exact dynamics. This problem is sometimes glossed over with the use of exponential propagators that are guaranteed to produce norm-conserving dynamics on a projected Hamiltonian. Some more sophisticated approaches adjust the configuration space in an attempt to mitigate




FIG. 7. The dynamics of a spin observable at a short time step, $dt = 0.01$, are indistinguishable when generated with equations of motion determined by the time embedded discrete variational principle (TEDVP) (dashed lines) and the McLachlan variational principle (MVP) (solid lines). Results are shown for propagations restricted to the space of single excitations from the initial state (S), double excitations (SD), and the full space (Ex). The associated error metrics, $\mathcal{N}_1(t)$ for the TEDVP (dashed) and $\mathcal{N}_2(t)$ for the MVP (solid), also yield nearly identical results, displaying peaks correlated with qualitative deviations from the exact trajectory.

the error [50].

This discrepancy is at the center of the difference between the approximate dynamics derived from the discrete variational principle here and the approximate dynamics derived from the McLachlan variational principle such as the multi-configurational time-dependent Hartree method. Mathematically, this results from the non-commutativity of the exponential map and projection operator defined above. That is $P_{\mathcal{B}_i(t)} \mathcal{T} \left(e^{-i \int dt' H(t')} \right) P_{\mathcal{B}_i(t)} \neq \mathcal{T} \left(\exp \left(-i \int dt' P_{\mathcal{B}_i(t')} H(t') P_{\mathcal{B}_i(t')} \right) \right)$ for a Hermitian operator H . In an approximate method derived from the McLachlan or any of the other differential time-dependent variational principles, the projection is performed on the Hamiltonian. As the projection of any Hermitian operator yields another Hermitian operator, the dynamics generated from the projection are guaranteed to be unitary if a sufficiently accurate exponential

propagator is used. Conversely, projection of a unitary operator, as prescribed by the TEDVP, does not always yield a unitary operator. Thus for an approximate basis, one expects norm conservation to be violated, where the degree of violation is related to the severity of the configuration space truncation error. This leads us to define a metric of truncation error which we term the instantaneous norm loss. We define this as

$$\mathcal{N}_1(t) = 1 - \|P_{\mathcal{B}_i} U_t P_{\mathcal{B}_i} |\tilde{\Psi}_t\rangle\|^2 \quad (35)$$

where $|\tilde{\Psi}_t\rangle$ is always assumed to be normalized, which in practice means that a renormalization is used after each time step here. However, as we proved above, in the limit of a short time step, with dynamics generated by a Hamiltonian, the TEDVP must contain essentially the same content as the McLachlan variational principle. For this reason, we propose an additional metric which is given by

$$\mathcal{N}_2(t) = \|(H(t) - P_{\mathcal{B}_i} H(t) P_{\mathcal{B}_i}) |\tilde{\Psi}_t\rangle\|^2 dt^2 \quad (36)$$

$$= \|V(t) |\tilde{\Psi}_t\rangle\|^2 dt^2 \quad (37)$$

Where $H(t)$ is the physical Hamiltonian. This is motivated by appealing to the McLachlan variational principle and substituting from the exact Schrödinger equation that $i\partial_t = H |\Psi_t\rangle$ where H is the full (non-projected) Hamiltonian. By defining $V(t) = (H(t) - P_{\mathcal{B}_i} H(t) P_{\mathcal{B}_i})$, we recognize this as a perturbation theory estimate of the error resulting from the configuration basis truncation at a given point in time.

To examine the quality of these metrics and to better understand the consequences of the non-commutativity of the exponential map and projection, we return to the sample spin system. In this case, we choose a basis for the propagation space based entirely on the initial state, and do not allow it to change dynamically in time as before. We perform simulations in the space of single excitations (S) from the initial state, double excitations (SD), and in the full Hilbert space (Ex). Dynamics from the TEDVP are generated by first building the exact propagator then projecting to the desired basis set while dynamics from the McLachlan variational principle (MVP) are modeled by projecting the Hamiltonian into the target basis set and exponentiating. A renormalization is used after each time step in the first case. Although one could perform the simulation with a timestep where timestep error is negligible, we remove this component of the calculation for this example by making the Hamiltonian time-independent. This allows direct analysis of the effect of timestep on non-commutativity deviations.

In Fig. 7 we show the dynamics for the two approximate truncation levels (S and SD) with both methods and their associated error metrics ($\mathcal{N}_1(t)$ and $\mathcal{N}_2(t)$). Deviations in the qualitative features of the observable occur after even the first peak of the proposed metrics. In this particular simulation, the configuration interaction

with singles and doubles spans all but one state in the full Hilbert space. The lack of this one state results in the large qualitative errors present, associated with the first and subsequent peaks present in these error metrics. The impact of later peaks is more difficult to discern, due to error in the wavefunctions, which accumulates as the propagation proceeds.

As predicted by the connection between the TEDVP and the McLachlan variational principle, while \mathcal{N}_1 and \mathcal{N}_2 are not identical for each case, in the short time limit they yield extremely similar information, which is highlighted in Fig. 7 displaying the resulting longer time dynamics for a time step of $dt = 0.01$. In Fig. 8, however we explore the effects of a significantly larger time step and begin to discern the result of the non-commutativity discussed here. Recalling that because the Hamiltonian is time-independent in this case, the propagator used is numerically exact in both instances, so this effect is not a result of what would be traditionally called time step error, resulting from intrinsic errors of an integrator. Interestingly, it is observed that $\mathcal{N}_1(t)$ begins to decay to a nearly constant value. This occurs because the action of projection after exponentiation breaks the degeneracy of the spectrum of the unitary operator, resulting in eigenvalues with norms different than 1. As a result, repeated action and renormalization of the operator is analogous to a power method for finding the eigenvector associated with the largest eigenvalue. This effect is exacerbated by taking long time steps.

Conclusions

In this manuscript, we introduce a new variational principle for time-dependent dynamics inspired by the Feynman clock originally employed for quantum computation. Unlike other previously-proposed variational principles, the proposed TEDVP approach involves the solution of an eigenvalue problem for the entire time propagation. This perspective allows for readily employing many of the powerful truncation techniques from quantum chemistry and condensed-matter physics, that have been developed for the exact diagonalization problem. The possibility of parallelize “in time” by building a large eigenvalue problem is important, as there are not many practical methods for achieving this in large-scale supercomputers. We introduce two novel error metrics for the TEDVP that allow one to characterize the basis approximations involved. We demonstrated the features of the method by simulating the dynamics of a Hydrogen molecule and a molecular effective spin Hamiltonian. Further research directions include the use of other approximate techniques for the time dynamics such as the use of perturbation theory [51] or coupled-cluster approaches [52].




FIG. 8. The dynamics of a spin observable at a large time step, $dt = 0.5$, shows significant differences resulting from the non-commutativity of projection and exponentiation in dynamics generated by the time embedded discrete variational principle (TEDVP) (dashed lines) and the McLachlan variational principle (MVP) (solid lines). Results are shown for propagations restricted to the space of single excitations from the initial state (S), double excitations (SD), and the full space (Ex). The corresponding error metrics, $\mathcal{N}_1(t)$ for the TEDVP (dashed) and $\mathcal{N}_2(t)$ for the MVP (solid), differ considerably in this case.

Acknowledgments

We would like to acknowledge David Tempel for his valuable comments on the manuscript. J.M. is supported by the DOE Computational Science Graduate Fellowship under grant number DE-FG02-97ER25308. A.A.-G. receives support from the HRL Laboratories, LLC under award number M1144-201167-DS and UCSD under grant number FA9550-12-1-004. We also thank the Corning Foundation, the Camille and Henry Dreyfus Foundation, and the Alfred P. Sloan Foundation for their continuing support of this research.

Appendix: On the general construction of eigenvalue problems from dynamics

We have presented one path for constructing eigenvalue problems from quantum dynamics problems so far, however it is instructive to illuminate precisely which part of our procedure allowed this. To do this, we will slightly generalize the Floquet-type Hamiltonians and demonstrate that the time embedding was the crucial feature that allows one to recast a dynamics problem as an eigenvalue problem. This is in addition to the choice to work in an integrated framework, which we will show simply allows for a convenient choice of basis.

Recall the definition of a Floquet-type Hamiltonian given by

$$F(t) = H(t) - i\partial_t \quad (38)$$

If one considers a finite time evolution for a length of time T , this operator is Hermitian in the basis of Fourier functions states given by

$$|\Psi_{nj}(t)\rangle = |\Phi_j\rangle |n\rangle = |\Phi_j\rangle e^{2\pi i n t / T} \quad (39)$$

where $|\Phi_j\rangle$ is a time-independent state of the physical system when considering the generalized inner-product

$$\langle n' | \langle \Phi_i | \Phi_j \rangle | n \rangle = \frac{\langle \Phi_i | \Phi_j \rangle}{T} \int_0^T dt' e^{-2\pi i n' t' / T} e^{2\pi i n t' / T} \quad (40)$$

This is the the generalized Hilbert space first introduced by Sambe [53] and generalized by Howland [54]. Because this operator is Hermitian in this basis, by noting the similarity to the Lagrange Variational principle

$$\delta L = \delta \int_0^T dt \langle \Psi(t) | F | \Psi(t) \rangle = 0 \quad (41)$$

minimization of L on this linear basis of Fourier states yields a Hermitian eigenvalue problem. Thus the time evolution can be reconstructed by solving the full time-independent eigenvalue problem in this basis, or by constructing a surrogate evolution operator as in the (t, t') formalism of Peskin and Moiseyev [14]. The use of Fourier basis states to express time dependence is natural given the form of the operator F . That is, matrix elements of the derivative operator ∂_t have a trivial analytic expression in this basis. This represents the same general idea we have been discussing here, which is to consider states in a system-time Hilbert space. However, as the solution of this variational problem will may yield a stationary point rather than a true minimum [30], ground state techniques are not appropriate for this particular operator. Moreover, this operator is not in general Hermitian when considering arbitrary basis functions of time. For example arbitrary choices of plane waves not corresponding to the traditional Fourier basis will yield a non-Hermitian operator. The operator $F' = (1/2)(F + F^\dagger)$,

which has been used in the past for the construction of approximate dynamics [55], is Hermitian, however it still suffers from a pathology that the optimal solution represents a stationary point rather than a minimum. However, the operator

$$G = F^\dagger F = (H(t) - i\partial_t)^\dagger (H(t) - i\partial_t) \quad (42)$$

is always Hermitian and positive semi-definite by construction. Thus one can expand the system-time Hilbert space in any linear basis of time, and the optimal path in that space will be the ground state eigenvector of the operator G utilizing the above generalized inner-product, assuming we have broken degeneracy by introducing the correct initial state. This can be viewed as an application of the McLachlan time dependent variational principle. From this, we see it is the expression systems in a system-time Hilbert space which allows for the eigenvalue problem construction. Moreover, we note that this is not limited to the use of Fourier time basis states, and that many more expressions of time dependence may be utilized to construct an eigenvalue problem within this framework.

One may ask then, what was the objective of working in the an integrated formalism, defined using unitary operators rather than differential operators. To see this, consider evaluating the system at a point in time with the Fourier construction above. One has to expand the system in all time basis functions and evaluate them at a time t . When one usually considers time, however, they are thinking of a parametric construction where a number t simply labels a specific state of a system. Embedding into the system-time Hilbert space with this idea would be most naturally expressed as delta-functions. However matrix elements of ∂_t on this basis can be difficult to construct. In the ancillary time system framework used in the TEDVP, however, time is easily expressed as a discrete parametric variable. One might also consider the use of discretized derivatives in the operator G . However, balancing the errors in numerical derivatives and the increasing difficulty of solving the problem with the number of simultaneously stored time steps can be practically difficult.

-
- [1] R. Feynman, Int. J. Theor. Phys. **21**, 467 (1982).
 - [2] R. P. Feynman, Opt. News **11**, 11 (1985).
 - [3] B. A. Lippmann and J. Schwinger, Phys. Rev. **79**, 469 (1950).
 - [4] E. J. Heller, J. Chem. Phys. **64**, 63 (1976).
 - [5] A. Kerman and S. Koonin, Annals of Physics **100**, 332 (1976).
 - [6] R. Jackiw and A. Kerman, Phys. Lett. A **71**, 158 (1979).
 - [7] R. Balian and M. Vénéroni, Phys. Rev. Lett. **47**, 1353 (1981).

- [8] E. Deumens, A. Diz, R. Longo, and Y. Öhrn, *Rev. Mod. Phys.* **66**, 917 (1994).
- [9] J. A. Poulsen, *J. Chem. Phys.* **134**, 034118 (2011).
- [10] J. Haegeman, J. I. Cirac, T. J. Osborne, I. Pizorn, H. Verschelde, and F. Verstraete, *Phys. Rev. Lett.* **107**, 070601 (2011).
- [11] K. F. Milfeld and R. E. Wyatt, *Phys. Rev. A* **27**, 72 (1983).
- [12] S. H. Autler and C. H. Townes, *Phys. Rev.* **100**, 703 (1955).
- [13] D. R. Dion and J. O. Hirschfelder, “Time-dependent perturbation of a two-state quantum system by a sinusoidal field,” in *Adv. Chem. Phys.* (John Wiley and Sons, Inc., 2007) pp. 265–350.
- [14] U. Peskin and N. Moiseyev, *J. Chem. Phys.* **99**, 4590 (1993).
- [15] M. Nielsen and I. Chuang, *Quantum Computation and Quantum Information*, Cambridge Series on Information and the Natural Sciences (Cambridge University Press, 2000).
- [16] I. Kassal, J. D. Whitfield, A. Perdomo-Ortiz, M.-H. Yung, and A. Aspuru-Guzik, *Annu. Rev. Phys. Chem.* **62**, 185 (2011).
- [17] M.-H. Yung, J. D. Whitfield, S. Boixo, D. G. Tempel, and A. Aspuru-Guzik, (2012), arXiv:1203.1331 [quant-ph].
- [18] A. Aspuru-Guzik, A. D. Dutoi, P. J. Love, and M. Head-Gordon, *Science* **309**, 1704 (2005).
- [19] I. Kassal, S. P. Jordan, P. J. Love, M. Mohseni, and A. Aspuru-Guzik, *Proc. Natl. Acad. Sci. USA* **105**, 18681 (2008).
- [20] H. Wang, S. Kais, A. Aspuru-Guzik, and M. R. Hoffmann, *Phys. Chem. Chem. Phys.* **10**, 5388 (2008).
- [21] I. Kassal and A. Aspuru-Guzik, *J. Chem. Phys.* **131**, 224102 (2009).
- [22] B. P. Lanyon, J. D. Whitfield, G. G. Gillett, M. E. Goggin, M. P. Almeida, I. Kassal, J. D. Biamonte, M. Mohseni, B. J. Powell, M. Barbieri, A. Aspuru-Guzik, and A. G. White, *Nat. Chem.* **2**, 106 (2010).
- [23] Z. Li, H. Yung, Man-Hong Chen, D. Lu, J. D. Whitfield, X. Peng, A. Aspuru-Guzik, and J. Du, *Sci. Rep.* **1** (2011).
- [24] G. K.-L. Chan and M. Head-Gordon, *J. Chem. Phys.* **116**, 4462 (2002).
- [25] F. Verstraete, J. J. García-Ripoll, and J. I. Cirac, *Phys. Rev. Lett.* **93**, 207204 (2004).
- [26] M. M. Wilde, *From Classical to Quantum Shannon Theory* (2011) arXiv:1106.1445.
- [27] L. Amico, R. Fazio, A. Osterloh, and V. Vedral, *Rev. Mod. Phys.* **80**, 517 (2008).
- [28] P. A. M. Dirac, *Math. Proc. Cambridge* **26**, 376 (1930).
- [29] J. Frenkel, *Wave Mechanics* (Clarendon Press, Oxford, 1934).
- [30] A. McLachlan, *Mol. Phys.* **8**, 39 (1964).
- [31] J. C. Slater, *Phys. Rev.* **34**, 1293 (1929).
- [32] S. F. Boys, *Proc. R. Soc. Lond. A Math. Phys. Sci.* **200**, 542 (1950).
- [33] E. Farhi, J. Goldstone, S. Gutmann, and M. Sipser, (2000), arXiv:quant-ph/0001106.
- [34] A. Kitaev, A. Shen, M. Vyalii, and N. Vyalii, *Classical and Quantum Computation*, Graduate Studies in Mathematics (American Mathematical Society, 2002).
- [35] F. Coester and H. Kümmel, *Nucl. Phys.* **17**, 477 (1960).
- [36] J. Cizek, *J. Chem. Phys.* **45**, 4256 (1966).
- [37] R. J. Bartlett and M. Musiał, *Rev. Mod. Phys.* **79**, 291 (2007).
- [38] M. Beck, A. Jäckle, G. Worth, and H.-D. Meyer, *Phys. Rep.* **324**, 1 (2000).
- [39] J. Broeckhove, L. Lathouwers, E. Kesteloot, and P. V. Leuven, *Chem. Phys. Lett.* **149**, 547 (1988).
- [40] A. Mizel, *Phys. Rev. A* **70**, 012304 (2004).
- [41] J. Lions, Y. Maday, and G. Turinici, *Comptes Rendus de l’Academie des Sciences Series I Mathematics* **332**, 661 (2001).
- [42] L. Baffico, S. Bernard, Y. Maday, G. Turinici, and G. Zerah, *Phys. Rev. E* **66**, 057701 (2002).
- [43] A. Mizel, M. W. Mitchell, and M. L. Cohen, *Phys. Rev. A* **65**, 022315 (2002).
- [44] W. J. Hehre, R. F. Stewart, and J. A. Pople, *J. Chem. Phys.* **51**, 2657 (1969).
- [45] I. Kuprov, N. Wagner-Rundell, and P. Hore, *J. Magn. Reson.* **189**, 241 (2007).
- [46] H. J. Hogben, P. J. Hore, and I. Kuprov, *J. Chem. Phys.* **132**, 174101 (2010).
- [47] M. Luban, F. Borsa, S. Bud’ko, P. Canfield, S. Jun, J. K. Jung, P. Kögerler, D. Mentrup, A. Müller, R. Modler, D. Procissi, B. J. Suh, and M. Torikachvili, *Phys. Rev. B* **66**, 054407 (2002).
- [48] S. Habershon, *J. Chem. Phys.* **136**, 054109 (2012).
- [49] A. Castro, M. A. L. Marques, and A. Rubio, *J. Chem. Phys.* **121**, 3425 (2004).
- [50] T. Westermann and U. Manthe, *J. Chem. Phys.* **137**, 22A509 (2012).
- [51] T. Helgaker, P. Jorgensen, and J. Olsen, *Molecular Electronic Structure Theory* (Wiley, Sussex, 2002).
- [52] S. Kvaal, *J. Chem. Phys.* **136**, 194109 (2012).
- [53] H. Sambe, *Phys. Rev. A* **7**, 2203 (1973).
- [54] J. S. Howland, *Mathematische Annalen* **207**, 315 (1974).
- [55] D. V. Shalashilin and I. Burghardt, *J. Chem. Phys.* **129**, 084104 (2008).